

Pittsburg, Kansas Air Quality Monitoring Study



Mark J. Travers, PhD, MS
Department of Health Behavior
Roswell Park Cancer Institute

February 2010

Executive Summary


In October, 2009, indoor air quality was assessed in 9 locations; 2 smoke-free restaurants, and 7 bar/restaurants permitting indoor smoking, in Pittsburg Kansas.

The concentration of fine particle air pollution, PM_{2.5}, was measured with a TSI SidePak AM510 Personal Aerosol Monitor. PM_{2.5} is particulate matter in the air smaller than 2.5 microns in diameter. Particles of this size are released in significant amounts from burning cigarettes, are easily inhaled deep into the lungs, and cause a variety of adverse health effects including cardiovascular and respiratory morbidity and death.

Key findings of the study include:

- In the 7 locations permitting indoor smoking, there were, on average, 1.7 cigarettes burning during the visits. This translates to an average of 0.34 burning cigarettes per 100 cubic meters of air in these places.
- In the 7 locations permitting indoor smoking the level of fine particle air pollution was hazardous (PM_{2.5} = 291 µg/m³). This level of particle air pollution is 73 times higher than in the smoke-free restaurants in Pittsburg and 29 times higher than outdoor air in Kansas.
- Employees in the locations with indoor smoking are exposed to levels of air pollution 5 times higher than safe annual levels established by the U.S. Environmental Protection Agency due to their occupational exposure to tobacco smoke pollution.

Figure 1. Average Level of Indoor Air Pollution in Pittsburg Locations Sampled



* Used for comparison purposes. Based on the 2008 average PM_{2.5} level from the EPA monitoring sites in Kansas (<http://www.epa.gov/air/data/>). The color-coded EPA Air Quality Index is also shown to demonstrate the magnitude of the measured particle levels

Introduction

Secondhand smoke (SHS) contains at least 250 chemicals that are known to be toxic or carcinogenic, and is itself a known human carcinogen,[1] responsible for an estimated 3,000 lung cancer deaths annually in *never smokers* in the U.S., as well as more than 35,000 deaths annually from coronary heart disease in *never smokers*, and respiratory infections, asthma, Sudden Infant Death Syndrome, and other illnesses in children.[2] Although population-based data show declining SHS exposure in the U.S. overall, SHS exposure remains a major public health concern that is entirely preventable.[3, 4] Because requiring smoke-free environments is the most effective method for reducing SHS exposure in public places,[5] Healthy People 2010 Objective 27-13 encourages all states and the District of Columbia to establish and to enforce smoke-free air laws in public places and worksites.[6]

Currently in the U.S., 27 states, Washington D.C., and Puerto Rico have passed strong smoke-free air laws that include restaurants and bars. The states are Arizona, California, Colorado, Connecticut, Delaware, Hawaii, Illinois, Iowa, Maine, Maryland, Massachusetts, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oregon, Rhode Island, Utah, Vermont, Washington, and Wisconsin.¹ Well over 50% of the U.S. population is now protected from secondhand smoke in all public places.[7] Nine Canadian provinces and territories also have comprehensive smoke-free air laws in effect. Hundreds of cities and counties across the U.S. have also taken action, as have whole countries including Ireland, Scotland, Uruguay, Norway, New Zealand, Sweden, Italy, Spain, England and France.

The goal of this study was to determine the level of fine particle air pollution in Pittsburg, Kansas venues permitting smoking and to compare this to smoke-free venues and outdoor air. There is currently no local smoke-free air law in Pittsburg. There is also no state level smoke-free air law in Kansas, although Kansas law does not preempt the passage of local smoke-free air laws.

It is hypothesized that: 1) indoor particle air pollution levels will be significantly lower in locations that do not allow indoor smoking compared to locations where smoking is permitted; and, 2) across all venues sampled, the degree of indoor particle air pollution will be correlated with the amount of smoking.

¹ The Michigan law goes into effect 5/1/2010 and the Wisconsin law goes into effect 7/5/2010.

Methods

Overview

On October 23rd and 24th, 2009, indoor air quality was assessed in 9 locations; 2 smoke-free restaurants, and 7 bar/restaurants permitting indoor smoking, in Pittsburg Kansas.

Measurement Protocol

A minimum of 30 minutes was spent in each venue. The number of people inside the venue and the number of burning cigarettes were recorded every 15 minutes during sampling. These observations were averaged over the time inside the venue to determine the average number of people on the premises and the average number of burning cigarettes. Room dimensions were also determined using a combination of any or all of the following techniques; a sonic measuring device, counting of construction materials of a know size such as floor tiles, or estimation. Room volumes were calculated from these dimensions. The active smoker density was calculated by dividing the average number of burning cigarettes by the volume of the room in meters.

A TSI SidePak AM510 Personal Aerosol Monitor (TSI, Inc., St. Paul, MN) was used to sample and record the levels of respirable suspended particles in the air. The SidePak uses a built-in sampling pump to draw air through the device where the particulate matter in the air scatters the light from a laser. This portable light-scattering aerosol monitor was fitted with a 2.5 μm impactor in order to measure the concentration of particulate matter with a mass-median aerodynamic diameter less than or equal to 2.5 μm , or $\text{PM}_{2.5}$. Tobacco smoke particles are almost exclusively less than 2.5 μm with a mass-median diameter of 0.2 μm . [8] The Sidepak was used with a calibration factor setting of 0.32, suitable for secondhand smoke. [9, 10] In addition, the SidePak was zero-calibrated prior to each use by attaching a HEPA filter according to the manufacturer's specifications.

The equipment was set to a one-minute log interval, which averages the previous 60 one-second measurements. Sampling was discreet in order not to disturb the occupants' normal behavior. For each venue, the first and last minute of logged data were removed because they are averaged with outdoors and entryway air. The remaining data points were averaged to provide an average $\text{PM}_{2.5}$ concentration within the venue.

TSI SidePak AM510 Personal Aerosol Monitor



PM_{2.5} is the concentration of particulate matter in the air smaller than 2.5 microns in diameter. Particles of this size are released in significant amounts from burning cigarettes, are easily inhaled deep into the lungs, and are associated with pulmonary and cardiovascular disease and mortality.

Teams of trained testers from the Clean Air Pittsburg did the sampling and researchers from Roswell Park Cancer Institute analyzed the data.

Statistical Analyses

The primary goal was to assess the difference in the average levels of PM_{2.5} between smoke-free and smoking-permitted establishments. Statistical significance is assessed using an independent samples *t* test on the log-transformed PM_{2.5} concentrations. PM_{2.5} concentrations are log-transformed to achieve normality and homogeneity of variances. The second hypothesis was tested by using all 9 sample visits and correlating the average smoker densities to the PM_{2.5} levels using the Spearman rank correlation coefficient (r_s). Descriptive statistics including the venue volume, number of patrons, and average smoker density (i.e., number of burning cigarettes) per 100 m³ are reported for each venue and averaged for all venues.

Results

A summary of each location visited is shown in Table 1. The average PM_{2.5} level in the 7 locations permitting indoor smoking was 291 µg/m³, compared to 4 µg/m³ in the 2 smoke-free locations (Figure 1). The difference between PM_{2.5} levels in smoking and smoke-free places is large and statistically significant ($t(8) = 2.05$, $p=.04$, $r=0.61$).


Table 1. Fine Particle Air Pollution in Pittsburg Kansas Bars and Restaurants

Venue Number	Size (m ³)	Average # people	Average # burning cigs	Active smoker density*	Average PM _{2.5} level (µg/m ³)
No Observed Smoking					
1	1408	24	0.0	0.00	3
2	96	14	0.0	0.00	5
Average (n=2)	752	19	0.0	0.00	4
Smoking Observed					
3	297	10	0.5	0.17	181
4	233	39	1.0	0.43	58
5	339	11	2.8	0.83	438
6	1712	25	2.0	0.12	278
7	565	63	3.0	0.53	442
8	505	17	1.0	0.20	476
9	1269	17	1.3	0.10	163
Average (n=7)	703	26	1.7	0.34	291

*Average number of burning cigarettes per 100 cubic meters.

In the 7 smoking-permitted locations the average number of burning cigarettes was 1.7 which corresponds to an average smoker density (ASD) of 0.34 burning cigarettes per 100 m³. In contrast, the 2 smoke-free locations had an ASD of 0.00, as no burning cigarettes were observed in these locations. Looking at all 9 sample visits, PM_{2.5} level was significantly correlated with the active smoker density ($r_s=0.71$, $p=0.032$), indicating that the amount of indoor smoking is the primary driver of the indoor particle pollution levels.

Figure 1. Average Level of Indoor Air Pollution in Pittsburg Locations Sampled



* Used for comparison purposes. Based on the 2008 average PM_{2.5} level from the EPA monitoring sites in Kansas (<http://www.epa.gov/air/data/>). The color-coded EPA Air Quality Index is also shown to demonstrate the magnitude of the measured particle levels

The real-time plot showing the level of indoor air pollution in each venue sampled is presented in Figures 2 on page 11. The real-time PM_{2.5} plot reveals the following results: 1) low background levels are observed outdoors; 2) high levels of indoor air pollution are observed in the venues where smoking was permitted; and 3) peak exposure levels in some venues where smoking was permitted reached levels far in excess of the average recorded level.

Discussion

The EPA cited over 80 epidemiologic studies in creating a particulate air pollution standard in 1997.[11] The EPA has recently updated this standard and, in order to protect the public health, the EPA has set limits of $15 \mu\text{g}/\text{m}^3$ as the average annual level of $\text{PM}_{2.5}$ exposure and $35 \mu\text{g}/\text{m}^3$ for 24-hour exposure.[11] In order to compare the findings in this study with the annual EPA $\text{PM}_{2.5}$ exposure standard, it was assumed that a full-time employee in the locations sampled that allow smoking works 8 hours, 250 days a year, is exposed to $291 \mu\text{g}/\text{m}^3$ (the average level in the sites with smoking) on the job, and is exposed only to background particle levels of $10 \mu\text{g}/\text{m}^3$ during non-work times. For a full-time employee their average annual $\text{PM}_{2.5}$ exposure is $74 \mu\text{g}/\text{m}^3$. The EPA average annual $\text{PM}_{2.5}$ limit is exceeded by 5 times due to their occupational exposure. In contrast, workers in the smoke-free location are exposed to an average particle concentration of $4 \mu\text{g}/\text{m}^3$ and, for a full-time employee in these Pittsburg venues, the average annual exposure is $9 \mu\text{g}/\text{m}^3$, a safe level according to the EPA. Based on the latest scientific evidence, the EPA staff currently proposes even lower $\text{PM}_{2.5}$ standards to adequately protect the public health,[12] making the high $\text{PM}_{2.5}$ exposures of people in smoking environments even more alarming.

Previous studies have evaluated air quality by measuring the change in levels of respirable suspended particles (RSP) between smokefree venues and those that permit smoking. Ott et al. did a study of a single tavern in California and showed an 82% average decrease in RSP levels after smoking was prohibited by a city ordinance.[13] Repace studied 8 hospitality venues, including one casino, in Delaware before and after a statewide prohibition of smoking in these types of venues and found that about 90% of the fine particle pollution could be attributed to tobacco smoke.[14] Similarly, in a study of 22 hospitality venues in Western New York, Travers et al. found a 90% reduction in RSP levels in bars and restaurants, an 84% reduction in large recreation venues such as bingo halls and bowling alleys, and a 58% reduction even in locations where only SHS from an adjacent room was observed at baseline.[15] A cross-sectional study of 53 hospitality venues in 7 major cities across the U.S. showed 82% less indoor air pollution in the locations subject to smokefree air laws, even though compliance with the laws was less than 100%.[16]

Other studies have directly assessed the effects SHS exposure has on human health. Rapid improvements in the respiratory health of bartenders were seen after a state smokefree workplace law was implemented in California[17]. Smokefree legislation in Scotland was associated with significant early improvements in symptoms, lung function, and systemic inflammation of all bar workers, while asthmatic bar workers also showed reduced airway inflammation and improved quality of life.[18] Farrelly et al. also showed a significant decrease in both salivary cotinine concentrations and sensory symptoms in hospitality workers after New York State's smokefree law prohibited smoking in their worksites.[19] A meta-analysis of the 8 published studies looking at the effects of smokefree air policies on heart attack admissions yielded an estimate of an immediate 19% reduction in heart attack admissions associated with these laws.[20]

The effects of passive smoking on the cardiovascular system in terms of increased platelet aggregation, endothelial dysfunction, increased arterial stiffness, increased atherosclerosis, increased oxidative stress and decreased antioxidant defense, inflammation, decreased energy production in the heart muscle, and a decrease in the parasympathetic output to the heart, are often nearly as large (averaging 80% to 90%) as chronic active smoking. Even brief exposures to SHS, of minutes to hours, are associated with many of these cardiovascular effects. The effects of secondhand smoke are substantial and rapid, explaining the relatively large health risks associated with secondhand smoke exposure that have been reported in epidemiological studies.[21]

The hazardous health effects of exposure to second-hand smoke are now well-documented and established in various independent research studies and numerous international reports. The body of scientific evidence is overwhelming: there is no doubt within the international scientific community that second-hand smoke causes heart disease, lung cancer, nasal sinus cancer, sudden infant death syndrome (SIDS), asthma and middle ear infections in children and various other respiratory illnesses. There is also evidence suggesting second-hand smoke exposure is also causally associated with stroke, low birthweight, spontaneous abortion, negative effects on the development of cognition and behavior, exacerbation of cystic fibrosis, cervical cancer and breast cancer. The health effects of secondhand smoke exposure are detailed in recent reports by the California Environmental Protection Agency[22] and the U.S. Surgeon General[23].

Conclusions

This study demonstrates that employees and patrons in Pittsburg bars and restaurants allowing indoor smoking are exposed to harmful levels of indoor air pollution resulting from indoor smoking. A comprehensive smoke-free air policy that prohibits indoor smoking in all indoor places is the only proven means to eliminate this exposure to toxic tobacco smoke pollution. This type of policy will result in improved quality of life and health outcomes for Pittsburg workers and residents.

Acknowledgments

This study was funded by the Tobacco Free Kansas Coalition.

Special thanks go to volunteers from Clean Air Pittsburg who collected the data for this study.

Support for Roswell Park Cancer Institute researchers was provided by the Flight Attendant Medical Research Institute.

Roswell Park Cancer Institute (RPCI) is America's first cancer center founded in 1898 by Dr. Roswell Park. RPCI is the only upstate New York facility to hold the National Cancer Center designation of "comprehensive cancer center" and to serve as a member of the prestigious National Comprehensive Cancer Network.

Over its long history, Roswell Park Cancer Institute has made fundamental contributions to reducing the cancer burden and has successfully maintained an exemplary leadership role in setting the national standards for cancer care, research and education.


The campus spans 25 acres in downtown Buffalo and consists of 15 buildings with about one million square feet of space. A new hospital building, completed in 1998, houses a comprehensive diagnostic and treatment center. In addition, the Institute built a new medical research complex and renovated existing education and research space to support its future growth and expansion.

For more information about Roswell Park and cancer in general, please contact the Cancer Call Center at 1-877-ASK-RPCI (1-877-275-7724).



Kansas Air Quality Monitoring Study Pittsburg, October, 2009

Figure 2



References

1. National Toxicology Program, *9th Report on Carcinogens 2000*. 2000, U.S. Department of Health and Human Services, National Institute of Environmental Health Sciences: Research Triangle Park, NC.
2. CDC, *Annual smoking-attributable mortality, years of potential life lost, and economic costs - United States, 1995-1999*. MMWR, 2002. **51**(14): p. 300-320.
3. U.S. Department of Health and Human Services, *Second national report on human exposure to environmental chemicals*. 2003, US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Environmental Health: Atlanta, GA.
4. U.S. Department of Health and Human Services, *Reducing tobacco use: a report of the Surgeon General*. 2000, US Government Printing Office: Washington, DC.
5. Hopkins, D.P., et al., *Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke*. Am J Prev Med, 2001. **20**(2 Suppl): p. 16-66.
6. U.S. Department of Health and Human Services, *Healthy People 2010: Objectives for Improving Health (Volume 2)*. 2000, U.S. Government Printing Office: Washington, DC.
7. American Nonsmokers' Rights Foundation. *Summary of 100% Smokefree State Laws and Population Protected by 100% U.S. Smokefree Laws*. 2009 7/1/2009 [cited 2009 Jul 6]; Available from: <http://www.no-smoke.org/pdf/SummaryUSPopList.pdf>.
8. Klepeis, N.E., et al., *Determining Size-Specific Emission Factors for Environmental Tobacco Smoke Particles*. Aerosol Science and Technology, 2003. **37**: p. 780-790.
9. Klepeis, N.E., W.R. Ott, and P. Switzer, *Real-Time Measurement of Outdoor Tobacco Smoke Particles*. Journal of the Air & Waste Management Association, 2007. **57**: p. 522-534.
10. Travers, M.J., *Smoke-free air policy: changing what's in the air and in the body*, in *Social and Preventive Medicine [Dissertation]*. 2008, State University of New York at Buffalo: Buffalo.
11. U.S. Environmental Protection Agency, *National ambient air quality standards for particulate matter; final rule*. Federal Register, 1997. **62**(138): p. 38651-38701.
12. U.S. Environmental Protection Agency. *Draft Staff Paper for Particulate Matter Fact Sheet*. 2005 [cited 2006 October 4]; Available from: http://www.epa.gov/airlinks/pdfs/pmstaff2_fact.pdf.
13. Ott, W., P. Switzer, and J. Robinson, *Particle concentrations inside a tavern before and after prohibition of smoking: evaluating the performance of an indoor air quality model*. J Air Waste Manag Assoc, 1996. **46**(12): p. 1120-1134.
14. Repace, J.L., *Respirable particles and carcinogens in the air of Delaware hospitality venues before and after a smoking ban*. J Occup Environ Med, 2004. **46**(9): p. 887-905.
15. Travers, M.J., et al., *Indoor Air Quality in Hospitality Venues Before and After the Implementation of a Clean Indoor Air Law-Western New York, 2003*. Morbidity and Mortality Weekly Report (MMWR), 2004. **53**(44): p. 1038-1041.
16. Travers, M.J., A. Hyland, and J.L. Repace, *7-City Air Monitoring Study (7-CAMS), March-April 2004*. 2004, Roswell Park Cancer Institute: Buffalo.
17. Eisner, M.D., A.K. Smith, and P.D. Blanc, *Bartenders' respiratory health after establishment of smoke-free bars and taverns*. JAMA, 1998. **280**(22): p. 1909-14.
18. Menzies, D., et al., *Respiratory symptoms, pulmonary function, and markers of inflammation among bar workers before and after a legislative ban on smoking in public places*. JAMA, 2006. **296**(14): p. 1742-8.
19. Farrelly, M.C., et al., *Changes in hospitality workers' exposure to secondhand smoke following the implementation of New York's smoke-free law*. Tob Control, 2005. **14**(4): p. 236-41.
20. Glantz, S.A., *Meta-analysis of the effects of smokefree laws on acute myocardial infarction: An update*. Preventive Medicine, 2008. **47**(4): p. 452-453.
21. Barnoya, J. and S.A. Glantz, *Cardiovascular effects of secondhand smoke: nearly as large as smoking*. Circulation, 2005. **111**(20): p. 2684-98.
22. California Environmental Protection Agency, *Proposed Identification of Environmental Tobacco Smoke as a Toxic Air Contaminant*. 2005, California Environmental Protection Agency, Air Resources Board, Office of Environmental Health Hazard Assessment.
23. U.S. Department of Health and Human Services, *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*. 2006, U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health: Atlanta, GA.